Journal of Surgery and Insights



Research Article

Enow Orock G, et al. J Surg Insights: JSI-100022

Gastrointestinal Pathologies in a Center with Limited Resources: Pilot Study at Bafoussam Regional Hospital, Cameroon

Enow Orock G^{1, 2*}, Enow Orock A², Ngounou E², Konang N Jl², Noubom M³, Nzeukou E¹, Tabeu Nm¹ and Luma H⁴

¹Department of Pathological Anatomy and Cytology, Bafoussam Regional Hospital, Cameroon

²Department of Biomedical Sciences, University of Buea, Cameroon

²Department of Gastroenterology, Bafoussam Regional Hospital, Cameroon

³Department of Medical Biology, Bafoussam Regional Hospital, Cameroon

⁴Department of Gastroenterology, General Hospital Douala, Cameroon

*Corresponding author: Enow Orock G, Department of Pathological Anatomy and Cytology, Bafoussam Regional Hospital, Cameroon, Tel: 237677716045; Email: enowrock24@yahoo.com

Citation: Enow Orock G, Enow Orock A, Ngounou E, Konang N Jl, Noubom M, et al. (2020) Gastrointestinal Pathologies in a Center with Limited Resources: Pilot Study at Bafoussam Regional Hospital, Cameroon. J Surg Insights: JSI-100022

Received date: 23 July, 2020; Accepted date: 27 July, 2020; Published date: 03 August, 2020

Introduction

Aim: Pathology of the digestive tract is common in Cameroon. Despite this situation, studies of gastrointestinal pathology are rare. The pathological anatomy service is almost non-existent in the Cameroonian suburbs. Bafoussam in the west of the country is a semi-urban area and one of the rare ones to have a pathological anatomy department. This is why we have decided to present the epidemiological data of the pathologies of the digestive tract of thislocality.

Méthods: We conducted a three-month prospective study (October 2019-January 2020), in the pathological anatomy department of the Bafoussam Regional Hospital in Cameroon with the aim of presenting the structural pathologies of the digestive tract listed in this institution.

Results: The pathology of the digestive tract has an incidence of approximately 400 in this locality, and represents 45.24% of all the pathologies identified in the pathological anatomy service of the Bafoussam Regional Hospital The age of our patients varies between 17 and 98 years old, with an average of 52.8. Women are slightly more represented (sex ratio M: F of 1: 1.3). Samples from the digestive tract represent 45% of all histological samples recorded in this department. The most common location, in decreasing order, is the stomach (84%), followed by the esophagus (5%), the colon and the rectum (4% each). On the other hand, other sites including the duodenum are rather rare. Of 100 digestive tract samples examined, 98 were pathological with a preponderance of inflammatory lesions (80.61%) against 19.39% of neoplastic nature. The inflammatory pathology was distributed uniformly in all age groups while the malignant pathologies were not observed until after 50 years. Malignant pathologies were more common in men (sex ratio M: F of 3: 1). Gastritis was the most common diagnosis (76.53%) and 94.66% of these gastritis were associated with Helicobacter pylori.

Conclusion: The pathology of the digestive tract is important in our community and dominated by the inflammatory type. The most common site is the stomach, most of the damage to which is caused by Helicobacter pylori. It would be desirable to actively seek out and eradicate Helicobacter pylori infection to decrease the incidence of digestive pathology. Further studies are recommended to establish the true epidemiologic trend of gastrointestinal pathology in our community.



Keywords: Bafoussam; Cameroon; Digestive; Pathology

Introduction

The gastrointestinal tract (IGT) is one of the most complex devices in the body because it has to perform several functions, including digestion and absorption of nutrients. Exposed to a continuous insult from the local macrobiota, the latter' sepithelium is constantly renewed [1]. Several pathological conditions can thus affect the TGI.

Digestive diseases affect millions of patients around the world with a significant economic impact including high cost of the healthcare system, absenteeism at work, and poor quality of life in patients [2-4].

Studies report that gastrointestinal pathology is common in Cameroon [5]. Another indicates that these pathologies constitute 10% of the problems of patients in consultation in the regional hospital of Limbe and at the regional hospital of Buea, in the southwest region of Cameroon [6]. Studies on histologically confirmed structural gastrointestinal pathology are rare in Cameroon and little done in tropical environments [7,8]. This scarcity is attributed to a lack of anatomy-pathological services in Cameroon. This work presents an overview of structural pathologies of the gastrointestinal tract in our community.

Materials and Methods

We conducted a prospective study over a period of 03 months (October 2019 - January 2020). The study concerned surgical specimens sent to the pathological anatomy laboratory, involving the gastrointestinal tract. We used the routine histological technique: samples fixed with 10% formalin, sections made with a microtome in 0.5mm section and then stained with haemateineosin. Finally, the slides are interpreted on a simple light microscope by an anatomopathologist. All TGI biopsies received during the study period are listed, assembled and analyzed. The results are presented in tabular form and graphs.

Results

Out of 221 histological samples received during the study period 100 concerned the digestive system (45.24%) (Figure I and II) and (Table I-VI).



Figure I : Distribution of patients by age.

The average age of the patients is 52.8 years (17-98 years). The most represented patients were those aged 60 and over (43.0%).



Figure II : Distribution of patients by sex.

There is a slight predominance of women in our series, with an M: F sex ratio of 1: 1.3.

Site	Number of samples	Percentage (%)
Esophagus	5	5.0
Stomach	84	84.0
Duodenum	2	2.0
Côlon	4	4.0
Rectum	4	4.0
Pancréas	1	1.0
Total	100	100.00%

Table I : Distribution of samples by anatomical site.

In our series, the most frequent localization of IGT disorders is gastric at 84%, and the rest distributed as follows: Esophagus (5%), Rectum (4%), Colon (4%), Duodenum (2%) and Pancreas (1%).

Pathological diagnosis	Workforce	Percentage	
Adenocarcinoma	8	8.0%	
Carcinoma	2	2.0%	
Duodenitis	1	1.0%	
Gastritis	75	75.0%	
Lymphoma	2	2.0%	
Esophagitis	1	1.0%	
Benign polyp	7	7.0%	
RCUH*	2	2.00%	
Without lesion	2	2.00%	
Total	100	100.00%	
Note:*RCUH: ulcerative colitis.			

 Table II: Distribution of samples according to pathological diagnosis.

Out of 98 lesions recorded, 80.61% of lesions are inflammatory in nature against 19.39% of neoplastic lesions. Gastritis is the most common diagnosis (76.53%).

Age range	0 - 19	20 - 29	30 - 39	40 - 49	50 - 59	60 - 69	≥70	Total
Pathological dg	-	-	-	-	-	-	-	-
Adenocarcinoma	-	-	-	-	3	5	-	8
Carcinoma	-	-	-	-	1	-	1	2
Duodenitis	-	1	-	-	-	-	-	1
Gastritis	3	10	12	11	10	13	16	75
Lymphoma	-	-	-	-	-	2	-	2
Esophagitis	-	-	-	-	-	-	1	1
Benign polyp	-	-	-	1	1	4	1	7
RCUH	-	-	-	2	-	-	-	2
Without lesion	-	1	1	-	-	-	-	2
Total	3	12	13	14	15	24	19	100

Table III: Pathological diagnosis by age group.

Inflammatory gastrointestinal pathology is distributed uniformly across all age groups, while malignant pathologies appear after 50 years.

Pathological diagnosis	F	М	Total
Adenocarcinoma	1	7	8
Carcinoma	1	1	2
Duodenitis	-	1	1
Gastritis	48	27	75
Lymphoma	1	1	2
Esophagitis	1	-	1
Benign polyp	3	4	7
Proctocolitis	-	2	2
Without lesion	-	2	2
Total	55	45	100

Table IV: Diagnostic pathology by Sex.

Gastrointestinal malignancies are more frequent than men (20.93%) that women (3.63%) have a sex ratio of M: F of 3: 1.

Site / Dg pathological	Workforce	Percentage
Colon	4	4.00%
Adenocarcinoma	1	1.00%
Benign polyp	1	1.00%
RCUH	1	1.00%
Without lesion	1	1.00%
Duodenum	2	2.00%
Duodenitis	1	1.00%

Benign polyp	1	1.00%
Stomach	84	84.00%
Adenocarcinoma	5	5.00%
Gastritis	75	75.00%
Lymphoma	2	2.00%
Benign polyp	1	1.00%
Without lesion	1	1,00%
Esophagus	5	5.00%
Carcinoma	2	2.00%
Esophagitis	1	1.00%
Benign polyp	2	2.00%
Pancreas	1	1.00%
Adenocarcinoma	1	1.00%
Rectum	4	4.00%
Adenocarcinoma	1	1,00%
Benign polyp	2	2.00%
RCUH	1	1.00%
Total	100	100.00%

Table V: Pathological diagnosis according to anatomical sites.

The stomach is the most represented site, recording 84% of samples from the gastrointestinal tract, and recording 84.6% of all lesions in our study.

Pathological diagnosis	Workforce	Percentage
Adénocarcinome	5	5.95%
Gastritis	75	89.29%
Without atrophy	43	51.19%
Without H.p	3	3.57%
With H.p	40	47.62%
With atrophy	32	38.10%
Without H.p	1	1.19%
With H.p	31	36.90%
Lymphoma	2	2.38%
Benign polyp	1	1.19%
Without lesion	1	1.19%
Total	84	100.00%

Table VI : Stomach lesions.

The lesions of the stomach are 89.29% gastritis; associated with atrophy (42.66%) and very often associated with Helicobacter pylori (94.66% of cases).

Discussion

The average age of our patients is 52.8 years (17-98 years). The most represented patients were those aged 60 and over (43.0%); see figure I. Our series is similar to the literature reported [9], in which the average age of patients was 55 years (23-80 years) with a higher frequency for the 40-60 year age group, unlike to that of some authors [10] whose mean age was 38.5 years (7-85 years), and 57% of patients with an age between 20 and 39 years. We note a slight predominance of women in our series, with an M: F sex ratio of 9:11 (Figure II), as in previous literature [11] whose M: F sex ratio was 107: 113, unlike another [10] which was predominantly male (sex ratio M: F of 707: 514).

Our study shows that the stomach is the most represented site, it stores 84% of samples from the digestive system see **Table** I. This trend is corroborated with the study by Nezha, which had a predominantly 40.74% gastric sample [9]. On the other hand, certain literature reports that digestive affections are very frequent in Africa with a strong predominance of colonic pathology [12,13].

80.61% of lesions in ourseries are inflammatory in nature see **Table II**. This percentage is very high compared to that found by Ndjitoyap et al, which was 54.65% [14]. This high frequency of inflammatory pathology in our context could be related to the multiple aggressive factors for the gastric mucosa which are part of the local diet of this population [15-18]. Indeed, of all the lesions observed in our study, gastritis is the most frequent diagnosis (76.53%). This frequency is higher than that reported with a predominance of gastritis at 27.79%. However, an ATTIA study in Ivory Coast reported twice as many duodenitis as gastritis [12].

An antecedent report finds malignant digestive diseases as the main cause of pathologies of this system with high mortality among all neoplasms, contributing to approximately 3 million deaths in 2012 worldwide. This report presents as the main digestive malignancies in descending order, hepatocellularcarcinoma (HCC), colorectal cancer, gastric cancer (GC), pancreatic cancer, cancer of the esophagus and oral cancer [18].

In our study we find a predominance of malignant lesions of the digestive tract in male patients with an M: F ratio of 3: 1 see **Table IV**. This opinion was reported by [19]. However, its incidence is characterized by a significant geographic disparity [20].

The inflammatory gastrointestinal pathology is distributed uniformly in all age groups while the malignant pathologies appear after 50 years see **Table III**. Malignant gastrointestinal lesions are more common in men (20.93%), with an M: F sex ratio of 3: 1 see **Table IV**. It seems clear to us that age is a risk factor for cancers of the gastrointestinal tract [16]. Indeed our result corroborates with some literature which reports that cancers of the gastrointestinal tract are rare before age 50 with a male predominance (sex ratio of 5: 2), and their incidence increases with age [16].

The stomach is the most affected site of TGI, it recorded 84.6% of all lesions in our study see **Table V**. This percentage is very high compared to that of [10], who had a predominance of gastric lesions at 44%.

In the lesions of the stomach are gastritis at 89.29% see **Table VI** and often associated with atrophy (42.66%) and very often associated with Helicobacter pylori (94.66% of cases), Helicobacter pylori being a major risk factor for gastric cancer according to several authors [21-23].

Conclusion

About 45.24% of the histological samples sent to the pathological anatomy department concern the digestive system. The inflammatory lesions of this system represent 80.61% of which 19.39% are neoplastic lesions. The cause of the predominance of inflammatory lesions remains to be determined. The most common diagnosis is gastritis, most often associated with Helicobacter pylori (94.66%). The eradication of Helicobacter pylori is very important in the strategy to combat digestive pathology.

Références

- Barker N, van EsJH, Kuipers J,Kujala P, van den Born M, etal. (2007) Identification of stem cells in small intestine and colon by marker gene Lgr5. Nature 449:1003-1007.
- Tougas G, Chen Y, Hwang P, Liu MM, Eggleston A (1999) Prevalence and impact of upper gastrointestinal symptoms in the Canadian population: findings from the DIGEST study. Domestic/International Gastroenterology Surveillance Study. Am J Gastroenterol 94:2845-2854.
- Drossman DA and Dumitrascu DL (2006) Rome III: New standard for functional gastrointestinal disorders. J Gastrointestin Liver Dis 15:237-241.
- Camilleri M and Williams DE (2000) Economic burden of irritable bowel syndrome. Proposed strategies to control expenditures. Pharmacoeconomics17:331-338.
- George EnowOrock G,Boniface Fogw E, Frank Zoun A, Alain Chicho M (2016)Epidemiolocal and Anatomo-Clinical Aspects of Ulcerative Colitis in a Referral Center in Cameroon and Review of Literature.Int. J. InnovativeHealthcare Res 4:1-8.
- EnowOrock GE, Fogwe BM, Nkounlack DC, Weledji EP, Chichom MA, et al. (2014) Aclinicopatologic study of digestive orgqn lesions in a low resource setting and review of literature. The case of Regional Hospital of Buea, Cameroon (Central Africa) AJIH 4: 4-13.
- Attia Y, Soubeyrand J, Beda Y, Gaudet D, Manlan K, et al. (1981) Place of colonicpathology in the Africanenvironment.Med. Afr. Noire 28: 349-351.
- Peghini M, Barabe P, Seurat P, Phillippon G, Morcillo R, et al. (1987) Recto-colicpolyps in Senegal, results of 1500 low endoscopies performed at the main hospital of DAKAR Med. Trop47: 361-364.
- Taoufiq N, Naim A, Bouchbika Z, Benchekroune N, Jouhadi H, et al. (2017)Gastro-Intestinal StromalTumors "GIST": current situation and news throughourexperiencerelating to 54 cases and a literaturereview. Pan Afr Med J 27: 165.
- Bougouma A, Drabo YJ, Serme AK, Zoungrana SL, Sombie AR, et al. (2002) Analysis of the results of 1221 hospital colonoscopies in Burkina Faso, 1994-1997. Bulletin de la Société de pathologie exotique 95: 50-52.
- 11. Djibril MA, M'Ba KB, Kaaga YL, Bagny A, EdouKA, et al. (2009) Upper digestive endoscopy in rural Africa in Togo ; Med Trop 69: 48-50.





- Camara BM, Manlan K, Attia Y (1985) The rarity of diverticulardisease in an Africanenvironment in the IvoryCoast. About 8 observations and the analysis of 1890 bariumenemaresults at the UniversityHospital of Cocody. Ann Univ Nation CI série B (Médecine) 19: 179-192.
- NditoyapNdam EC, Tzeuton C, Mbakop A, Pouepene J, Guemne TA, et al. (1990) Endoscopie digestive haute au Cameroun. Med AFr Noire 37: 453-456.
- 14. Potet F (1988) Histopathologie du tube digestif. Masson, Paris. Pp : 324.
- 15. FaivrJ (1998) Report presented to the 100th French Congress of Surgery. Epidemiology of stomach cancer, Paris. Pp :1-12.
- Attia Y, Gaudet D, Manlan K, Carsuzaa M (1981) Assessment of 15 months of upper digestive endoscopy at CocodyUniversityHospital. Rev. Med. Côte d'Ivoire 54: 16-22.
- Wada T, Fujiwara H, Morita SH, Fukagaw T, wet Katai H (2017) Incidence of and risk factors for preoperative deep venous thrombosis in patients undergoing gastric cancer surgery.Gastric Cancer20: 872-877.
- DaniyalM,AhmadS,AhmadM,AsifHM,AkramM,et al. (2015) Risk-FactorsandEpidemiologyofGastricCancerinPakistan. AsianPac J CancerPrev16:4821-4824.

- Melloukil,LazarN,BenyachouB,AqodadN,IbrahimiA (2014) Epidemiology of gastriccancer:Experience of a Moroccanhospital center. PanAfriMedJ 17:42.
- FormanD,NewellDG,FullertonF,YarnelIJW,StaceyAR,et al. (1991) Association between infection with Helicobacter pylori and risk of gastric Cancer: evidence from a prospective investigation.Br.Med.J 302:1302-1305.
- Nomura A, Stemmermann GN, Chyou P, Kato I, Perez-Perez GI, et al. (1991) Helicobacter pylori infection and gastric carcinoma in a population of Japanese- Americans in Hawaii. N. Engl. J. Med 325:1132-1136.
- Parsonnet J, Friedman GD, Vandersteen DP, Chang Y, VogelmanJH, et al. (1991) Helicobacter pylori infection and the risk of gastric carcinoma. N. Eng. J. Med 325:1127-1113.
- Talley NJ, Zinsmeister AR, Dimagno EP, Weaver A, Carpenter HA, et al. (1991) Gastric adenocarcinoma and Helicobacter pylori Infection. J. Nati. Cancer Inst83:1734-1739.